

A discriminant criterion of irreducibility

Evelia R. García Barroso and Janusz Gwoździewicz

November 19, 2009

Abstract

In this paper we give a criterion of irreducibility for a complex power series in two variables, using the notion of jacobian Newton diagrams, defined with respect to any direction. Moreover we study the singularity at infinity of a plane affine curve with one point at infinity for which the global counterpart of our main result holds.

1 Introduction

In [Gar-Gw] we give criteria of irreducibility for a complex power series in two variables, using the notion of jacobian Newton diagrams, defined with respect to a generic direction. In this paper we generalize these criteria to any direction and we use this new general criterion to study the branches at infinity of polynomial curves. The paper is organized as follows:

In 1.1 we recall the notion of the *Newton diagram*. Then in 1.2 we explain what we mean by the *discriminant curve* of an analytic mapping $F : (\mathbf{C}^2, 0) \rightarrow (\mathbf{C}^2, 0)$. If $D(u, v) = 0$ is an equation of the discriminant curve then the Newton diagram of D will be called *jacobian Newton diagram* of F and denoted $\mathcal{N}_J(F)$. At the end of the section we present formulas for computing equations of discriminants.

In Section 2 we consider $\mathcal{N}_J(l, f)$ where l is a regular function and f is a singular irreducible series. We shall call such diagrams the *Merle type diagrams*. We recall Merle's result that equisingularity class of f and the intersection multiplicity $(f, l)_0$ determine and are determined by $\mathcal{N}_J(l, f)$. In Theorem 2.3 we give necessary and sufficient conditions of arithmetical nature for a Newton diagram to be a Merle type diagram.

The main result of the paper is Theorem 3.1. It states that f is an irreducible power series if and only if $\mathcal{N}_J(l, f)$ is a Merle type diagram. We apply our irreducibility criterion to power series taken from Kuo's paper [Kuo].

2000 *Mathematics Subject Classification*: Primary 32S55; Secondary 14H20.

Key words and phrases: irreducible plane curve, jacobian Newton polygon, discriminant curve, branches at infinity.

The first-named author was partially supported by the Spanish Project PNMTM 2007-64007.

Finally we study the singularity at infinity of a plane affine curve with one point at infinity for which the global counterpart of our main result holds. This is interesting in the context of the Jacobian conjecture. Recall that Abhyankar proved in [Abh1] that this conjecture is settled affirmatively in the case where there is only one branch at infinity.

1.1 Newton diagrams of plane analytic curves

In this section we recall the notion of a Newton diagram and establish the notation. Write $\mathbf{R}_+ = \{x \in \mathbf{R} : x \geq 0\}$.

Let $f \in \mathbf{C}\{x, y\}$, $f(x, y) = \sum a_{i,j} x^i y^j$ be a non-zero convergent power series. Put $\text{supp } f := \{(i, j) : a_{i,j} \neq 0\}$. Then by definition the *Newton diagram* Δ_f , in the coordinates (x, y) , of f is

$$\Delta_f = \text{Convex Hull}(\text{supp } f + \mathbf{R}_+^2).$$

The basic property of Newton diagrams is that the Newton diagram of a product is the Minkowski sum of Newton diagrams. There is $\Delta_{fg} = \Delta_f + \Delta_g$ where $\Delta_f + \Delta_g = \{a + b : a \in \Delta_f, b \in \Delta_g\}$. In particular if f and g differ by an invertible factor $u \in \mathbf{C}\{x, y\}$, $u(0, 0) \neq 0$ then $\Delta_f = \Delta_g$. Thus the Newton diagram of a plane analytic curve is well defined because an equation of an analytic curve is given up to invertible factor.

After Teissier [Te2] we introduce *elementary Newton diagrams*. For $m, n > 0$ we put $\{\frac{n}{m}\} = \Delta_{x^n + y^m}$. We put also $\{\frac{n}{\infty}\} = \Delta_{x^n}$ and $\{\frac{\infty}{m}\} = \Delta_{y^m}$.

One can check that every Newton diagram $\Delta \subsetneq \mathbf{R}_+^2$ has a unique representation $\Delta = \sum_{i=1}^r \{\frac{L_i}{M_i}\}$ where *inclinations* of successive elementary diagrams form an increasing sequence (by definition the inclination of $\{\frac{L}{M}\}$ is L/M with the conventions that $L/\infty = 0$ and $\infty/M = +\infty$). We shall call this representation the *canonical form* of Δ .

Finally a Newton diagram is *convenient* if it intersects both coordinate axes.

1.2 Discriminant curve

Let $F = (p, q) : (\mathbf{C}^2, 0) \rightarrow (\mathbf{C}^2, 0)$ be an analytic mapping given by $(u, v) = (p(x, y), q(x, y))$ and such that $F^{-1}(0, 0) = \{(0, 0)\}$. Let $\text{jac}(p, q) = 0$ be the equation of the critical locus of F where $\text{jac}(p, q) = \frac{\partial p}{\partial x} \frac{\partial q}{\partial y} - \frac{\partial p}{\partial y} \frac{\partial q}{\partial x}$ is the usual jacobian determinant. The direct image of $\text{jac}(p, q) = 0$ by F is called the *discriminant curve* of F (see [Ca2]).

Assume that $D(u, v) = 0$ is an equation of the discriminant curve. Then Δ_D , in the coordinates (u, v) , is called the *jacobian Newton diagram* of F (see [Te3]). We will write $\mathcal{N}_J(p, q)$ for the jacobian Newton diagram.

Below we give some formulas for jacobian Newton diagrams and discriminant curves.

Formula 1.1 (Teissier's formula [Te1]) Assume that $\text{jac}(p, q) = h_1 \cdots h_r$ where h_i are irreducible series for $1 \leq i \leq r$. Then

$$\mathcal{N}_J(p, q) = \sum_{i=1}^r \left\{ \frac{(q, h_i)_0}{(p, h_i)_0} \right\}$$

where $(f, g)_0$ denotes the intersection number of f and g .

From now on we will only consider the mappings

$$(l, f) : (\mathbf{C}^2, 0) \rightarrow (\mathbf{C}^2, 0) \quad (1)$$

where l is a regular function (i.e. $l = ax + by + \text{higher order terms}$, $ax + by \neq 0$) and f is a singular series. Recall that a power series is called *singular* if its order is larger than one. Under these assumptions $\text{jac}(l, f) = 0$ is called the *polar curve of f with respect to l* . The inclinations of the elementary diagrams of the jacobian Newton diagram $\mathcal{N}_J(l, f)$ are called *polar quotients*. These notions were studied by many authors (see for example [Me] and [Eph] for irreducible case, and [Ca1], [Del1], [Del2], [Eg], [GB], [Gw-P12], [Ma1], [Ma2] and [Wall] among others for the reduced case). Also [Gw-Len-P1] is a survey of recent results. If the curves $l = 0$ and $f = 0$ are transverse then $\mathcal{N}_J(l, f)$ depends only on the equisingularity class of $f = 0$ (see [Te1]). Otherwise the jacobian Newton diagram may depend on relative position of curves $l = 0$ and $f = 0$ as the following example shows.

Example 1.2 Let $f = y^2 - x^5$ and let $l_1 = x$, $l_2 = y$, $l_3 = y - x^2$. Then $\text{jac}(l_1, f) = 2y$, $\text{jac}(l_2, f) = 5x^4$ and $\text{jac}(l_3, f) = x(5x^3 - 4y)$. By Teissier's formula

$$\begin{aligned} \mathcal{N}_J(l_1, f) &= \left\{ \frac{(f, y)_0}{(l_1, y)_0} \right\} = \left\{ \frac{5}{1} \right\} \\ \mathcal{N}_J(l_2, f) &= 4 \left\{ \frac{(f, x)_0}{(l_2, x)_0} \right\} = 4 \left\{ \frac{2}{1} \right\} = \left\{ \frac{8}{4} \right\} \\ \mathcal{N}_J(l_3, f) &= \left\{ \frac{(f, x)_0}{(l_3, x)_0} \right\} + \left\{ \frac{(f, 5x^3 - 4y)_0}{(l_3, 5x^3 - 4y)_0} \right\} = \left\{ \frac{2}{1} \right\} + \left\{ \frac{5}{2} \right\}. \end{aligned}$$

For any local analytic diffeomorphism $\Phi : (\mathbf{C}^2, 0) \rightarrow (\mathbf{C}^2, 0)$ the substitution $(l_1, f_1) = (l \circ \Phi, f \circ \Phi)$ does not affect the equation of the discriminant curve. Hence without loss of generality we may assume that $l = x$ (take such a Φ that $l \circ \Phi = x$).

Formula 1.3 If $f(x, y)$ is a convergent power series such that $f(0, y) = y^n + \text{higher order terms}$ and $\frac{\partial f}{\partial y}(x, y) = \text{unit} \prod_{j=1}^{n-1} [y - \gamma_j(x)]$ is a Newton-Puiseux factorization of $\frac{\partial f}{\partial y}$ then the discriminant of the mapping $(x, f) : (\mathbf{C}^2, 0) \rightarrow (\mathbf{C}^2, 0)$ has, up to an invertible factor, an equation

$$D(u, v) = \prod_{j=1}^{n-1} [v - f(u, \gamma_j(u))]. \quad (2)$$

See the Appendix for the proof.

Formula 1.4 *If $f(x, y) = y^n + a_1(x)y^{n-1} + \dots + a_n(x) \in \mathbf{C}\{x\}[y]$ is a Weierstrass polynomial, i.e. $a_i(0) = 0$ for every $i \in \{1, \dots, n\}$, then the discriminant of the mapping $(x, f) : (\mathbf{C}^2, 0) \rightarrow (\mathbf{C}^2, 0)$ has an equation $D(u, v) = 0$ where*

$$D(u, v) = \text{Discr}_y(f(u, y) - v) \quad (3)$$

is the classical discriminant of a polynomial in one variable y .

Proof. The discriminant $\text{Discr}_y(f(u, y) - v)$ is, up to an integer constant, equal to the resultant of polynomials $f(u, y) - v$ and $\frac{\partial f}{\partial y}(u, y)$. By the classical formula (see Theorem 10.10, Chapter I, [Walk]) the resultant of polynomials $P, Q \in K[Y]$, $Q = \prod_{i=1}^s (Y - \beta_i)$ where K is a field is, up to a sign, a product $\prod_{i=1}^s P(\beta_i)$. Hence under notations of Formula 1.3 $\text{Discr}_y(f(u, y) - v) = c \prod_{j=1}^{n-1} [f(u, \gamma_j(u)) - v]$ where c is a nonzero constant and Formula 1.4 follows. ■

Formula 1.5 *Let $f(x, y) = y^N + a_1(x)y^{N-1} + \dots + a_N(x) \in \mathbf{C}\{x\}[y]$. Assume that all nonzero roots of the polynomial $f(0, y)$ are simple. Then the discriminant of the mapping $(x, f) : (\mathbf{C}^2, 0) \rightarrow (\mathbf{C}^2, 0)$ is given by formula (3).*

Proof. Let $\frac{\partial f}{\partial y}(x, y) = N \prod_{i=1}^{N-1} (y - \gamma_i(x))$ be the Puiseux factorization of y -partial derivative. Take $\gamma_k(x)$ such that $\gamma_k(0) \neq 0$. Since $\gamma_k(0)$ is a root of $\frac{\partial f}{\partial y}(0, y)$ and all nonzero roots of $f(0, y)$ are simple there is $f(0, \gamma_k(0)) \neq 0$.

We get

$$\begin{aligned} \text{Discr}_y(f(u, y) - v) &= \text{const} \prod_{i=1}^{N-1} [v - f(u, \gamma_i(u))] \\ &= \text{const} \prod_{\gamma_i(0) \neq 0} [v - f(u, \gamma_i(u))] \prod_{\gamma_i(0)=0} [v - f(u, \gamma_i(u))] \\ &= \text{unit} \prod_{\gamma_i(0)=0} [v - f(u, \gamma_i(u))] \end{aligned}$$

which is, up to a unit, (1.3). ■

2 Jacobian Newton diagrams of irreducible series

In this section we consider mappings $(l, f) : (\mathbf{C}^2, 0) \rightarrow (\mathbf{C}^2, 0)$ under additional assumption that f is an irreducible singular power series. Then the curve $f = 0$ is often called a *plane singular branch*.

Consider

$$S(f) = \{(f, g)_0 : g \in \mathbf{C}\{x, y\} \text{ and } f \text{ does not divide } g\}.$$

Clearly $0 \in S(f)$ (take $g = 1$) and if $a, b \in S(f)$ then $a + b \in S(f)$ since the intersection number is additive, so $S(f)$ is a semigroup, called the *semigroup of the branch $f = 0$* .

For any regular curve $l = 0$ the semigroup $S(f)$ has the $(f, l)_0$ -minimal system of generators $\bar{b}_0, \bar{b}_1, \dots, \bar{b}_h$ defined by conditions

- (i) $\bar{b}_0 = (f, l)_0$,
- (ii) $\bar{b}_k = \min(S(f) \setminus (\mathbf{N}\bar{b}_0 + \dots + \mathbf{N}\bar{b}_{k-1}))$,
- (iii) $S(f) = \mathbf{N}\bar{b}_0 + \dots + \mathbf{N}\bar{b}_h$.

The sequence of generators can be characterized in purely arithmetical terms. Let us recall (see [Bre], [Za] for the generic case ($\bar{b}_0 = \text{ord} f$) and [Gw-Pl1] for the case when the curves $f = 0, l = 0$ are tangent)

Theorem 2.1 *Let $\bar{b}_0, \bar{b}_1, \dots, \bar{b}_h$ be a sequence of positive integers. Set $n_k = \gcd(\bar{b}_0, \dots, \bar{b}_{k-1}) / \gcd(\bar{b}_0, \dots, \bar{b}_k)$ for $k \in \{1, \dots, h\}$. Then the following two conditions are equivalent.*

- (i) *There is a singular branch $f = 0$ and a regular curve $l = 0$ such that $\bar{b}_0, \bar{b}_1, \dots, \bar{b}_h$ is the $(f, l)_0$ -minimal system of generators of the semigroup $S(f)$,*
- (ii) *the sequence $\bar{b}_0, \bar{b}_1, \dots, \bar{b}_h$ satisfies the conditions:*
 - (Z₁) $n_k > 1$ for $k \in \{1, \dots, h\}$ and $n_1 \cdots n_h = \bar{b}_0$,
 - (Z₂) $n_k \bar{b}_k < \bar{b}_{k+1}$ for $k \in \{1, \dots, h-1\}$.

Now we can state the result due to [Sm], [Me] and [Eph].

Theorem 2.2 (Smith–Merle–Ephraim) *Suppose that $f = 0$ is a singular branch and $l = 0$ is a regular curve. Let $\bar{b}_0, \dots, \bar{b}_h$ be the $(f, l)_0$ -minimal system of generators of the semigroup $S(f)$. Then with the notation introduced above*

$$\mathcal{N}_J(l, f) = \sum_{k=1}^h \left\{ \frac{(n_k - 1)\bar{b}_k}{(n_k - 1)n_1 \cdots n_{k-1}} \right\}. \quad (4)$$

If $\bar{b}_0, \dots, \bar{b}_h$ is the sequence satisfying the conditions (Z₁) and (Z₂) of Theorem 2.1 then we will write $\mathcal{M}(\bar{b}_0, \dots, \bar{b}_h)$ for the Newton diagram (4) and we call it the *Merle type diagram*. Let us note that the Newton diagram in formula (4) is written in the canonical form because quotients of the inclinations of successive elementary Newton diagrams, which are $\bar{b}_{k+1}/(n_k \bar{b}_k)$, are greater than 1 by Theorem 2.1.

Let us look at Example 1.2 in the light of Theorem 2.2. The curve $f = 0$ has the semigroup $S(f) = \mathbf{N}2 + \mathbf{N}5$. There is $(f, l_1)_0 = 2$, $(f, l_2)_0 = 5$, $(f, l_3)_0 = 4$ and is easy to verify that $\mathcal{N}_J(l_1, f) = \mathcal{M}(2, 5)$, $\mathcal{N}_J(l_2, f) = \mathcal{M}(5, 2)$ and $\mathcal{N}_J(l_3, f) = \mathcal{M}(4, 2, 5)$.

Theorem 2.3 Let $\Delta = \sum_{i=1}^h \{ \frac{L_i}{M_i} \}$ be a convenient Newton diagram written in its canonical form. Put $H_0 = 1$, $H_i = 1 + M_1 + \dots + M_i$ for $i \in \{1, \dots, h\}$ and $C_0 = H_h$, $C_i = H_{i-1}L_i/M_i$ for $i \in \{1, \dots, h\}$. Then Δ is a Merle type diagram if and only if the arithmetic conditions (i)–(iii) are satisfied

- (i) the quotients H_i/H_{i-1} are integers for $i \in \{2, \dots, h\}$,
- (ii) the quotients C_i are integers for $i \in \{1, \dots, h\}$,
- (iii) $\gcd(C_0, \dots, C_i) = C_0/H_i$ for $i \in \{1, \dots, h\}$.

Moreover in such a case $\Delta = \mathcal{M}(C_0, \dots, C_h)$.

Proof. Assume that Δ is a Merle type diagram $\mathcal{M}(\bar{b}_0, \dots, \bar{b}_h)$. Then $L_i = (n_i - 1)\bar{b}_i$ and $M_i = (n_i - 1)n_1 \dots n_{i-1}$ for $i \in \{1, \dots, h\}$. We have the equality $H_i = n_1 \dots n_i$. Indeed $H_1 = 1 + M_1 = 1 + (n_1 - 1) = n_1$ and $H_{i+1} = H_i + M_{i+1} = n_1 \dots n_i + (n_{i+1} - 1)n_1 \dots n_i = n_1 \dots n_{i+1}$ by the inductive hypothesis. It follows that $H_i/H_{i-1} = n_i$ hence condition (i) is satisfied.

It also follows that $C_i = \frac{H_{i-1}L_i}{M_i} = \frac{n_1 \dots n_{i-1}(n_i - 1)\bar{b}_i}{(n_i - 1)n_1 \dots n_{i-1}} = \bar{b}_i$. Hence condition (ii) is also satisfied.

It follows directly from the definition of the sequence n_i that $\gcd(\bar{b}_0, \dots, \bar{b}_i) = \bar{b}_0/(n_1 \dots n_i)$ for $1 \leq i \leq h$. Moreover by condition (Z₁) of Theorem 2.1 there is $n_1 \dots n_h = \bar{b}_0$ which gives $C_0 = H_h = \bar{b}_0$. Thus by condition (Z₁) of Theorem 2.1 $\gcd(C_0, \dots, C_i) = C_0/H_i$ for $i \in \{1, \dots, h\}$.

Now assume that conditions (i)–(iii) hold true for the Newton diagram Δ . We will show that the sequence $C_0, \dots, C_h$ satisfies arithmetical conditions of Theorem 2.1. It follows from (iii) that $n_i := \frac{\gcd(C_0, \dots, C_{i-1})}{\gcd(C_0, \dots, C_i)} = H_i/H_{i-1}$ for $i \in \{1, \dots, h\}$. Thus $n_i > 1$ for $i \in \{1, \dots, h\}$ and $n_1 \dots n_h = C_0$.

Because Δ is written in canonical form there is $L_i/M_i < L_{i+1}/M_{i+1}$ for $i \in \{1, \dots, h-1\}$. Multiplying these inequalities by $n_i H_{i-1} = H_i$ we get $n_i H_{i-1} L_i/M_i < H_i L_{i+1}/M_{i+1}$ which is equivalent with $n_i C_i < C_{i+1}$ for $i \in \{1, \dots, h-1\}$.

We checked that the sequence $C_0, \dots, C_h$ satisfies conditions (Z₁) and (Z₂) of Theorem 2.1. Moreover looking at the first part of the proof it is easy to see that $\Delta = \mathcal{M}(C_0, \dots, C_h)$. ■

3 Discriminant criterion of irreducibility

Theorem 3.1 Let $f = 0$ be a plane singular curve and let $l = 0$ be a regular curve. Then f is irreducible if and only if $\mathcal{N}_J(l, f)$ is a Merle type diagram. Moreover if $\mathcal{N}_J(l, f) = \mathcal{M}(\bar{b}_0, \dots, \bar{b}_h)$ then $f = 0$ has the semigroup $S(f) = \mathbb{N}\bar{b}_0 + \dots + \mathbb{N}\bar{b}_h$.

Example 3.2 Let $l = x$ and $f = y^n - x^m$. Then by (2) $D(u, v) = (v + u^m)^{n-1}$, hence $\mathcal{N}_J(x, f) = \left\{ \frac{(n-1)m}{n-1} \right\}$. Under notation of Theorem 2.3 there is $C_0 = H_1 = n$, $C_1 = m$ and conditions (i) and (ii) of Theorem 2.3 are clearly satisfied. Condition (iii) reduces to $\gcd(m, n) = 1$ and it is well-known that the curve $y^n - x^m = 0$ is irreducible if and only if m and n are co-prime.

The following two examples are taken from [Kuo] (see also [Abh2]).

Example 3.3 Let $f = (y^2 - x^3)^2 - x^7$. Then $\text{jac}(x, f) = 4y(y^2 - x^3) = 4y(y - x^{3/2})(y + x^{3/2})$. By Formula 1.3 we get $D(u, v) = (v - u^6 + u^7)(v + u^7)^2$. Hence $\mathcal{N}_J(x, f) = \left\{ \frac{6}{1} \right\} + \left\{ \frac{14}{2} \right\}$. Under notation of Theorem 2.3 there is $H_1 = 1 + 1 = 2$, $C_0 = H_2 = 1 + 1 + 2 = 4$, $C_1 = 6/1 = 6$, $C_2 = H_1 \cdot 14/2 = 14$ and because $\gcd(C_0, C_1, C_2) = 2 \neq 1$, it follows that $\mathcal{N}_J(x, f)$ is not a Merle type diagram. Therefore f is not irreducible.

Example 3.4 Let $f(x, y) = (y^2 - x^3)^2 - x^5y$. By Formula 1.4 $D(u, v) = -256v^3 + 256u^6v^2 + 288u^{13}v - 256u^{19} - 27u^{20}$ (we computed the discriminant using Sage) and the Newton diagram of the discriminant is $\mathcal{N}_J(x, f) = \left\{ \frac{6}{1} \right\} + \left\{ \frac{13}{2} \right\}$. It is easy to check that $\mathcal{N}_J(x, f)$ is a Merle type diagram $\mathcal{M}(4, 6, 13)$. Therefore f is irreducible with semigroup $S(f) = \mathbf{N}4 + \mathbf{N}6 + \mathbf{N}13$.

Example 3.5 Let $f(x, y) = x^8 + (x^2 + y^3)^3$. The jacobian Newton diagram of (x, f) is $\mathcal{N}_J(x, f) = \left\{ \frac{12}{2} \right\} + \left\{ \frac{48}{6} \right\}$ which is not a Merle type diagram. Note that in this example $x = 0$ is not transverse to $f(x, y) = 0$.

Corollary 3.6 Let $f(x, y) = y^N + a_1(x)y^{N-1} + \dots + a_N(x) \in \mathbf{C}[x, y]$. Assume that the curve $f(x, y) = 0$ intersects $x = 0$ only at the point $(0, y_0)$. Then the curve $f(x, y) = 0$ is analytically irreducible at $(0, y_0)$ if and only if the Newton diagram of $\text{Discr}_y(f(u, y) - v)$ is a Merle type diagram.

Proof. Put $\tilde{f}(x, y) = f(x, y + y_0)$. Then $f(x, y) = 0$ is analytically irreducible at $(0, y_0)$ if and only if $\tilde{f}(x, y) = 0$ is analytically irreducible at $(0, 0)$. Since $\text{Discr}_y(f(u, y) - v) = \text{Discr}_y(\tilde{f}(u, y) - v)$ the result follows from Formula 1.5. ■

4 Proof of Theorem 3.1

The proof is based on Theorem 1 of [Gar-Gw]:

Theorem 4.1 Let $f, g \in \mathbf{C}\{x, y\}$ be such that $\mathcal{N}_J(x, f) = \mathcal{N}_J(x, g)$. Assume that $x = 0$ is transverse to the curves $f = 0$ and $g = 0$. If f is irreducible then g is also irreducible.

and on the following lemma

Lemma 4.2 Let f where $f(0, y) = y^n + \text{higher order terms}$ be a convergent power series and let N be a positive integer. Put $\tilde{f}(x, y) = f(x^N, y)$. Then

- (i) if N and n are coprime integers then f is irreducible if and only if $\tilde{f}$ is irreducible,
- (ii) if $N > n$ then $\tilde{f} = 0$ is transverse to $x = 0$,
- (iii) $\mathcal{N}_J(x, \tilde{f}) = L(\mathcal{N}_J(x, f))$ where $L : \mathbf{R}^2 \rightarrow \mathbf{R}^2$ is a linear automorphism given by $L(i, j) = (Ni, j)$.

Proof.

Proof of (i). Assume that $f = f_1 f_2$. Then $\tilde{f}(x, y) = f_1(x^N, y) f_2(x^N, y)$. It follows that if $\tilde{f}$ is irreducible then f is irreducible.

In order to show the implication in opposite direction assume that f is irreducible. Recall (see Theorem 2.1, Chapter IV, [Walk]) that the curve $f = 0$ where $\text{ord} f(0, y) = n$ is a branch if and only if there exists a convergent power series $\phi(t)$ such that $f(t^n, \phi(t)) = 0$ and the greatest common divisor of the set $\{n\} \cup \text{supp } \phi$ equals 1.

Let $\phi(t)$ be such a series and let $\tilde{\phi}(t) = \phi(t^N)$. Then $\tilde{f}(t^N, \tilde{\phi}(t)) = f(t^{nN}, \phi(t^N)) = 0$ and since n and N are co-prime the greatest common divisor of the set $\{n\} \cup \text{supp } \tilde{\phi} = \{n\} \cup N \cdot \text{supp } \phi$ equals 1. Consequently $\tilde{f} = 0$ is a branch.

Proof of (ii). By the assumption $N > n$ the homogeneous initial part of the series $f(x^N, y)$ is y^n . This gives (ii).

Proof of (iii). Let $\frac{\partial f}{\partial y}(x, y) = \text{unit} \prod_{j=1}^{n-1} [y - \gamma_j(x)]$ be the Newton-Puiseux factorization of $\frac{\partial f}{\partial y}$. By Formula 1.3 the discriminant of the mapping (x, f) has an equation $D(u, v) = \prod_{j=1}^{n-1} [v - f(u, \gamma_j(u))]$. Because $\frac{\partial \tilde{f}}{\partial y}(x, y) = \frac{\partial f}{\partial y}(x^N, y)$ there is $\frac{\partial \tilde{f}}{\partial y}(x, y) = \text{unit} \prod_{j=1}^{n-1} [y - \gamma_j(x^N)]$ and consequently the discriminant of the mapping $(x, \tilde{f})$ has an equation $\tilde{D}(u, v) = \prod_{j=1}^{n-1} [v - \tilde{f}(u, \gamma_j(u^N))] = \prod_{j=1}^{n-1} [v - f(u^N, \gamma_j(u^N))] = D(u^N, v)$. Comparing Δ_D with $\Delta_{\tilde{D}}$ we get (iii). ■

Now let us prove Theorem 3.1. Suppose that $\mathcal{N}_J(l, f) = \mathcal{N}_J(l, g)$ where g is an irreducible power series. Applying an analytic change of coordinates we may assume that $l = x$. Take an integer $N > 0$ such that conclusions of (i) and (ii) of Lemma 4.2 are satisfied for $\tilde{f}(x, y) = f(x^N, y)$ and $\tilde{g}(x, y) = g(x^N, y)$. It follows from (iii) of Lemma 4.2 that $\mathcal{N}_J(x, \tilde{f}) = \mathcal{N}_J(x, \tilde{g})$. Since $\tilde{f}$ and $\tilde{g}$ satisfy assumptions of Theorem 4.1, $\tilde{f}$ is an irreducible power series. Hence by (i) of Lemma 4.2 f is also irreducible.

5 Discriminant criterion of irreducibility at infinity

Let $p(x, y)$ be a complex polynomial of degree $n > 0$. Let $C \subset \mathbf{P}^2(\mathbf{C})$ be the projective closure of the curve $p(x, y) = 0$. Assume that C intersects the line at infinity at only one point Q . The purpose of this section is to give a criterion

for local analytical irreducibility of the curve C at Q without passing to local coordinates centered at Q . For this we need some notions.

Let $g(x, y)$ be a polynomial of positive degree such that $g(x, 0) \neq 0$ and $g(0, y) \neq 0$ (in other words its Newton diagram is convenient). Let $\mathcal{P}_0(g)$ be the boundary in $\mathbf{R}_+^2$ of Δ_g and $\mathcal{P}_\infty(g)$ be the boundary in $\mathbf{R}_+^2$ of $\Delta_\infty(g) = \text{Convex Hull}(\text{supp } g \cup \{(0, 0)\})$. We call these sets the *Newton polygon of g at zero* and the *Newton polygon of g at infinity* respectively.

Theorem 5.1 *Let $p(x, y)$ be a complex polynomial of degree $n > 0$ without multiple factors and let C be the projective closure of $p(x, y) = 0$. Assume that C intersects the line at infinity at only one point $Q \neq (0 : 1 : 0)$. Put $D_\infty(x, t) := \text{Disc}_y(p(x, y) - t)$ and let $L : \mathbf{Z}^2 \rightarrow \mathbf{Z}^2$ be the affine transformation defined by $L(i, k) = (n(n-1) - i - nk, k)$. Then the curve C is analytically irreducible at Q if and only if $L(\mathcal{P}_\infty(D_\infty))$ is the Newton polygon at zero of a Merle type diagram.*

Proof. Let $P(x, y, z) = z^n p(\frac{x}{z}, \frac{y}{z})$ be a homogeneous equation of the curve C . Assume that C intersects the line at infinity only at $Q = (1 : y_0 : 0)$. Then $p(x, y) = P(x, y, 1)$ and $f(y, z) := P(1, y, z) = 0$ is the affine equation of C in coordinates y, z . In these coordinates the point Q becomes $(y_0, 0)$. Since the curve C intersects the line $z = 0$ only at Q the polynomial $f(y, z)$ satisfies the assumptions of Corollary 3.6.

Put $D(x, z, t) = \text{Disc}_y(P(x, y, z) - t)$. We have that $D_\infty = D(x, 1, t)$ and $D_0 := \text{Disc}_y(f(z, y) - t) = D(1, z, t)$. Since $P(x, y, z)$ is a homogeneous polynomial of degree n , giving to the variable t the weight n , and the other variables the weight 1, the polynomial $D(x, z, t)$ is quasi-homogeneous of degree $n(n-1)$ (see Theorem 10.9, Chapter I, [Walk]). In particular any term $c_{ijk}x^i z^j t^k$ of $D(x, z, t)$ corresponds with the point (i, j, k) of the hyperplane $\Pi \equiv i + j + nk = n(n-1)$. Moreover such point determines the term $c_{ijk}x^i t^k$ of D_∞ and the term $c_{ijk}z^j t^k$ of D_0 . Put $L : \mathbf{Z}^2 \rightarrow \mathbf{Z}^2$ defined by $L(i, k) = (n(n-1) - i - nk, k)$. Then $\text{supp } D_0 = L(\text{supp } D_\infty)$. The last relation gives $\mathcal{P}_0(D_0) = L(\mathcal{P}_\infty(D_\infty))$. By Corollary 3.6 the curve C is analytically irreducible at Q if and only if the Newton diagram Δ_{D_0} is a Merle type diagram. ■

Remark 5.2 *Let $p(x, y)$ be a polynomial of degree n which has one point at infinity different from $(0 : 1 : 0)$. Let us denote by q the maximal inclination of $\mathcal{P}_0(D_0)$. After [Pl], the Abhyankar-Moh inequality (see [A-M]) is equivalent to $q < n$. Note also that the Abhyankar-Moh inequality is equivalent to equisingularity at infinity of the family $p(x, y) - t = 0$. After [Kra] this is also equivalent to the statement that all segments of $\mathcal{P}_\infty(D_\infty)$ have positive slopes.*

Example 5.3 *Let $p(x, y) = x + (x + y^3)^3$ be a polynomial in $\mathbf{C}[x, y]$ which corresponds to the projective curve C defined by $P(x, y, z) = xz^8 + (xz^2 + y^3)^3 =$*

0. The only point at infinity of C is $Q = (1 : 0 : 0)$. Moreover $D_\infty = (x + x^3 - t)^2(x - t)^6$ and $\mathcal{P}_\infty(D_\infty)$ has only two segments joining the point $(0, 8)$ to $(6, 6)$ and this one to $(12, 0)$. The transformation of $\mathcal{P}_\infty(D_\infty)$ by $L(i, k) = (72 - i - 9k, k)$ is a polygon of two segments joining the point $(0, 8)$ to $(12, 6)$ and this one to $(60, 0)$. This polygon is the Newton polygon of $\Delta = \{\frac{12}{2}\} + \{\frac{48}{6}\}$. Since Δ is not a Merle type diagram, by Theorem 5.1 the curve C is not analytically irreducible at Q . Observe that the local equation $P(1, y, z) = 0$ of the curve C was studied in Example 3.5.

Appendix

The purpose of this section is the proof of Formula 1.3. We establish a more general statement. Using basic properties of direct image we show that if $F = (x, f)$ and $h(x, y) = \prod_{j=1}^m [y - \gamma_j(x)]$ is the Newton-Puiseux factorization of the power series h then the direct image $F_*(h = 0)$ has an equation

$$\prod_{j=1}^m [v - f(u, \gamma_j(u))] = 0. \quad (5)$$

Lemma 5.4 *Let $\phi(t) \in \mathbf{C}\{t\}$, $\phi(0) = 0$ be a convergent power series and let m be a positive integer. Let $V \subset (\mathbf{C}^2, 0)$ be the image of the mapping $(\mathbf{C}, 0) \ni t \rightarrow (t^m, \phi(t)) \in (\mathbf{C}^2, 0)$. Then V has an analytic (not necessarily reduced) equation $g = 0$ where*

$$g(x, y) = \prod_{\epsilon^m=1} [y - \phi(\epsilon x^{1/m})].$$

Lemma 5.4 is an easy corollary of:

Theorem 5.5 (Puiseux Theorem) *Let $h(x, y)$ be an irreducible power series and let $n = \text{ord} f(0, y) < \infty$. Then there exists $\phi(t) \in \mathbf{C}\{t\}$, $\phi(0) = 0$ such that $h(t^n, \phi(t)) = 0$. Moreover h , up to an invertible factor, is equal to*

$$H(x, y) := \prod_{\epsilon^n=1} [y - \phi(\epsilon x^{1/n})]. \quad (6)$$

Conversely, if $(\mathbf{C}, 0) \ni t \rightarrow (t^n, \phi(t)) \in (\mathbf{C}^2, 0)$ is an analytic parametrization of a plane branch then $H(x, y) = 0$ is its reduced equation.

Proof. (of Lemma 5.4) Let d be the greatest common divisor of $\{m\} \cup \text{supp } \phi$. Then there exists a power series ϕ_0 such that $\phi(t) = \phi_0(t^d)$.

Put $n = m/d$. Since the greatest common divisor of $\{n\} \cup \text{supp } \phi_0$ equals 1 the mapping $t \rightarrow (t^n, \phi_0(t))$ is an analytic parametrization of V . Let ϵ_m be the m -th primitive root of unity and let $\epsilon_n = \epsilon_m^d$. By Puiseux' Theorem V has an equation $h(x, y) = \prod_{i=1}^n [y - \phi_0(\epsilon_n^i x^{1/n})] = 0$.

Put $g = h^d$. Then $g(x, y) = \prod_{i=1}^n [y - \phi_0(\epsilon_n^i x^{1/n})]^d = \prod_{i=1}^{nd} [y - \phi_0(\epsilon_n^i x^{1/n})] = \prod_{i=1}^m [y - \phi_0((\epsilon_m^i x^{1/m})^d)] = \prod_{i=1}^m [y - \phi(\epsilon_m^i x^{1/m})]$ and we get Lemma 5.4. Observe that g is not reduced in the case $d > 1$. ■

Now we prove (5) under assumption that $h = 0$ is a branch.

Lemma 5.6 *Let $h(x, y) = \prod_{i=1}^n [y - \gamma_i(x)]$ be an irreducible power series. Then the direct image of the branch $h = 0$ by the mapping $(x, f) : (\mathbf{C}^2, 0) \rightarrow (\mathbf{C}^2, 0)$ has an equation $\prod_{i=1}^n [v - f(u, \gamma_i(u))] = 0$.*

Proof. By Puiseux' Theorem we may assume that $\gamma_i(x) = \phi(\epsilon_n^i x^{1/n})$ for $i = 1, \dots, n$ where $t \rightarrow (t^n, \phi(t))$ is an analytic parametrization of a branch $h = 0$ and ϵ_n is the n -th primitive root of unity.

Thus the image of the curve $h = 0$ by the mapping (x, f) is given by $(u, v) = (t^n, f(t^n, \phi(t)))$. By Lemma 5.4 it has an equation $g(u, v) = \prod_{i=1}^n [v - f(u, \phi(\epsilon_n^i u^{1/n}))] = \prod_{i=1}^n [v - f(u, \gamma_i(u))] = 0$. Moreover $(g, u)_0 = (h, x)_0 = n$ which shows, after Projection Formula (see page 64 of [Ca1]), that $g = 0$ is the direct image of the curve $h = 0$. ■

Because the equation of the direct image of a curve $h_1 h_2 = 0$ is the product of equations of direct images of $h_i = 0$ ($i = 1, 2$) formula (5) holds also in the case when h is a reducible power series.

References

- [Abh1] S.S. Abhyankar, *Lectures on expansion techniques in algebraic geometry*, Tata Institute of Fundamental Research, Bombay, 1977.
- [Abh2] S.S. Abhyankar, *Irreducibility Criterion for Germs of Analytic Functions of Two Complex Variables*, Advances in Mathematics 74, (1989), 190-257.
- [A-M] S.S. Abhyankar and T. Moh, *Embeddings of the line in the plane*, J. Reine Angew. Math. 276 (1975), 148-166.
- [Bre] H. Bresinsky, *Semigroups corresponding to algebroid branches in the plane*, Proc. of the AMS, vol. 32, No 2, (1972), 381-384.
- [Ca1] E. Casas-Alvero, *Singularities of plane curves*, London Mathematical Society, Lecture Note Series 276, 2000.
- [Ca2] E. Casas-Alvero, *Discriminant of a morphism and inverse images of plane curve singularities*, Math. Proc. Camb. Phil. Soc. (2003), 135, 385-394.
- [Del1] F. Delgado de la Mata, *An arithmetical factorization for the critical point set of some map germs from $\mathbf{C}^2$ to $\mathbf{C}^2$* , Singularities (Lille 1991), 61-100. London Math. Soc. Lecture Note Ser. 201, 1994.
- [Del2] F. Delgado de la Mata, *A factorization theorem for the polar of a curve with two branches*, Compositio Math. 92 (1994), 327-375.
- [Eg] H. Eggers, *Polarinvarianten und die Topologie von Kurvensingularitäten*, Bonner Mathematische Schriften 147, 1983.

- [Eph] R. Ephraim, *Special polars and curves with one place at infinity* (P. Orlik ed.), Proc. of Symp. in Pure Math., Vol 40 Part 1, AMS, Providence, 1983, 353–359.
- [GB] E. García Barroso, *Sur les courbes polaires d’une courbe plane réduite*, Proc. London Math. Soc. 81, Part 1, (2000), 1-28.
- [Gar-Gw] E. García Barroso, J. Gwoździewicz, *Characterization of jacobian Newton polygons of plane branches and new criteria of irreducibility*, arXiv:085.4257
- [Gw-Len-Pł] J. Gwoździewicz A. Lenarcik and A. Płoski, *Polar Invariants of Plane Curve Singularities: Intersection Theoretical Approach*, Demonstratio Math. (to appear).
- [Gw-Pł1] J. Gwoździewicz and A. Płoski, *On the Approximate Roots of polynomials*, Annales Polonici Mathematici LX3 (1995), 199–210.
- [Gw-Pł2] J. Gwoździewicz and A. Płoski, *On the polar quotients of an analytic plane curve*, Kodai Math. J. **25** (2002), 43-53.
- [Kra] T. Krasieński, *The level sets of polynomials in two variables and the Jacobian conjecture*, Acta Univ. Łódź Folia Math. (published dissertation) 1991 (in Polish).
- [Kuo] T.C. Kuo, *Generalized Newton-Puiseux Theory and Hensel’s Lemma in $\mathbb{C}[[x, y]]$* , Canadian Journal of Mathematics, Vol. 41, No 6, (1989), 1101-1116.
- [Ma1] H. Maugendre, *Discriminant d’un germe $(g, f) : (\mathbb{C}^2, 0) \longrightarrow (\mathbb{C}^2, 0)$ et quotients de contact dans la résolution of $f.g$* , Annales de la Faculté de Sciences de Toulouse, 6^e srie, tome 3, n 3, (1998), 497-525.
- [Ma2] H. Maugendre, *Discriminant of a germ $\phi : (\mathbb{C}^2, 0) \longrightarrow (\mathbb{C}^2, 0)$ and Seifert fibred manifolds*, Journal London Mathematical Society 1999 59 (1), 207-226.
- [Me] M. Merle, *Invariants polaires des courbes planes*, Invent. Math. 41, (1977) 103-111.
- [Pł] A. Płoski, *Polar quotients and singularities at infinity of polynomials in two complex variables*, Annales Polonici Mathematici LXXVIII.1 (2002), 49-58.
- [Sm] H. J. S. Smith, *On the higher singularities of plane curves*, Proc. London Math. Soc. 6 (1875), 153–182.
- [Tel] B. Teissier, *Variétés polaires.I. Invariants polaires des singularités des hypersurfaces*, Invent. Math. 40, (1977) 267-292.

- [Te2] B. Teissier, *The hunting of invariants in the geometry of discriminants*, Proc. Nordic summer school, 1976. Per Holm, editor, Sijthoff and Noordhoff 1978, p.565-677.
- [Te3] B. Teissier, B. Teissier, *Jacobian Newton polyhedra and equisingularity*, Proc. Kyoto Singularities Symposium, RIMS, 1978.
- [Walk] R. J. Walker, *Algebraic Curves*, Princeton University Press, New Jersey, 1950.
- [Wall] C.T.C. Wall, *Chains on the Eggers tree and polar curves*, Rev. Mat. Iberoamericana 19, 2, (2003) 745-754.
- [Za] O. Zariski, *Le problème des modules pour les branches planes*, Centre de Maths, Ecole Polytechnique, 1975. Reprinted by Hermann, Paris 1986.

Evelia Rosa García Barroso
 Departamento de Matemática Fundamental
 Facultad de Matemáticas, Universidad de La Laguna
 38271 La Laguna, Tenerife, España
 e-mail: ergarcia@ull.es

Janusz Gwoździwicz
 Department of Mathematics
 Technical University
 Al. 1000 L PP7
 25-314 Kielce, Poland
 e-mail: matjg@tu.kielce.pl